

# Teaching Multivariate Analysis to Business-Major Students

Wing-Keung Wong and Teck-Wong Soon - Kent Ridge, Singapore

## 1. Introduction

During the last two or three decades, multivariate statistical analysis has become increasingly popular. The theory has made great progress, and with the rapid advances in computer technology, routine applications of multivariate statistical methods are implemented in several statistical software packages, making it simple even for the novice to undertake fairly sophisticated multivariate statistical analysis of data at their disposal.

While this is certainly a welcome development, we find, on the other hand, that many users of statistical packages are unable to appreciate what they are doing, and this is particularly true for multivariate statistical methods. With the increasing use of multivariate statistical methods by business analysts, it is important for business-major students to develop an understanding of multivariate statistical methods. Even though business executives are not generally required to undertake sophisticated statistical analysis themselves, they are often presented with reports and articles based on such analysis. Furthermore, many executives have heard about these techniques and would like to use them as analytic and decision support tools.

The traditional approach to the teaching of multivariate statistical analysis, as exemplified by Anderson (1958), relies heavily on advanced matrix mathematics. On the other hand, Flury and Riedwyl (1988) suggest that it is possible to understand most of the basic ideas underlying multivariate statistical analysis without a mastery of such mathematics, provided that these are conveyed with the help of real data sets. Since most business data do not follow the usual normality assumption, there are often (possibly severe) limitations in the use of some of the standard multivariate statistical techniques. Real data sets are therefore required not only to illustrate the statistical techniques concerned, but also to clarify the assumptions needed for these techniques to be valid.

In this paper, we propose a *non-mathematical data-driven approach* for teaching multivariate statistical methods to business-major students. Despite this, we are mindful of the need for students to know some basic linear algebra and univariate statistical concepts. Such basic knowledge provides students with the foundation necessary for the application of the appropriate multivariate statistical procedures and for the interpretation of results.

## 2. Business data and the approach to analyse them

Business data sets are usually large and undifferentiated. They are therefore generally closer to data encountered in the social sciences and differ somewhat from the more precise data encountered in the physical and natural sciences. Large and undifferentiated data sets contain so many inter-relationships that it is virtually impossible to make sense of them without first arriving at a summary description. Multivariate statistical analysis, when applied to such data sets, should allow us to "explore" the data sets with a view to discovering, describing and understanding the major inter-relationships. While multivariate statistical techniques allow us to analyse, verify, test, and prove various hypotheses, it should be emphasised that with a large undifferentiated data set, we should, at least in the initial stages, be less involved with building a specific statistical model, or with the formal procedures of statistical inference. Thus, when teaching multivariate statistical analysis to business-major students, we should concentrate on the descriptive techniques rather than on the theory of the multivariate normal distribution, beginning with simple univariate statistical analysis leading to standard multivariate techniques such as principal component analysis, factor analysis, and cluster analysis which are applicable when the data is measured on a continuous or interval scale. Once the basic ideas have been conveyed, it would be a simple matter to introduce students to other related techniques, such as correspondence analysis and multi-dimensional scaling.

## 3. Illustrative example

To illustrate the approach outlined above, we consider the sample data set 'CARDATA' provided with the computer package STATGRAPHICS. The data set comprises 155 observations with 11 variables as follows: *mpg*, *cylinders*, *displace*, *horsepower*, *accel*, *year*, *weight*, *origin*, *make*, *model* and *price*. This data set is small enough for most computer packages to handle and large enough to approximate the type of data encountered in practice. Faced with such a data set, students are generally at a loss to know how to begin to analyse it, particularly when the objectives of the analysis have not been specified.

An obvious approach would be to begin by examining each of the variables in turn. Such an initial examination of the data has the merit of enabling students to obtain a better feel for the data. For example, on examining the variable *mpg*, we might consider the summary statistics tabulated in Table 1.

TABLE 1  
Summary statistics for *mpg*

Number of Observations	154
Mean	28.79
Standard Deviation	7.37
Coefficient of Variation	25.62

To enable students to familiarise themselves with the data, more elaborate univariate analysis can be undertaken. For example, students may construct histograms, plot stem-and-leaf diagrams, etc. Note also that because some variables may have missing values, the total number of observations reported may be less than 155.

Once the initial examination of data has begun, students might be led to raise further questions. For example, how does *mpg* vary with other variables? If the other variable in question is a classifying (i.e. nominal-scaled) variable, the observations can be grouped before analysis. As an illustration, consider the case when the other variable is *origin*. Table 2 presents the summary statistics.

TABLE 2  
Summary statistics for *mpg* by *origin*

	United States	Europe	Japan
Number of Observations	85	25	44
Mean	25.26	32.55	33.48
Standard Deviation	6.12	8.18	5.27
Coefficient of Variation	24.22	25.13	15.75

These summary statistics suggest that Japanese cars with the highest *mpg* are more economical than American or European cars. Students will then be led naturally to consider the possible reasons for this. Is it the case that Japanese cars are smaller or less powerful? Consideration of such questions will lead them to examine the variables *weight* and *horsepower*. Summary statistics presented in tables similar to Table 2 suggest that it is indeed the case that Japanese cars are lighter (i.e. smaller) and less powerful.


### 3.1 Regression and correlation analysis

The simple initial examination of data described above will prompt students to consider the relationship between two or more variables. They now know that Japanese cars are more fuel-economical because they are lighter and less powerful. Students will then be led to consider the relationship between fuel consumption, *mpg* and *weight* or *horsepower*. The suggestion of plotting a scatterplot of *mpg* against *weight* or *horsepower* can then be given. Students may then attempt to draw a "line of best fit" through the scatterplot. Figure 1, which shows the result of this exercise, indicates that *mpg* decreases as *weight* increases. Students could then be asked to consider how best to fit the line and thus led to "least squares estimation", obtaining the line of best fit to be:

$$mpg = 55.89 - 0.0101 \times weight$$

suggesting that for every lb increase in *weight*, the fuel consumption decreases by 0.01 *mpg*. Furthermore, students can be led to ask how much of the variation of *mpg* can be explained by the variable *weight*. This leads naturally to  $R^2$  which is the proportion of the variation of the variable *mpg* that is "explained" by *weight*. In this case,

$R^2 = 0.6874$ , suggesting that 68.74% of the variation may be explained by the linear regression.



Scatterplot of *mpg* vs *weight*

While discussing  $R^2$ , it can be explained that in the simple linear regression case (when there is only one explanatory variable),  $R^2$  is just the square of the sample linear correlation coefficient. In this way, students are introduced naturally to the fact that the correlation coefficient is a measurement of the closeness of the sample observations to a straight line. Once this idea is conveyed, students can discuss more complicated issues such as the testing of hypotheses about the parameters of the straight line as well as the assumptions needed to ensure that such tests are valid. Indeed, the appropriateness of the underlying assumptions of the linear regression model can also be examined by careful study of the scatter diagram which would lead students to consider residual analysis and issues such as normality, outliers, etc.

### 3.2 *Extension to multiple regression analysis*

The detailed consideration of a simple linear regression analysis such as that discussed above extends naturally to multiple linear regression analysis. While students may experiment fitting a regression of *mpg* on *horsepower* or *accel*, they will soon find this rather *ad hoc* and would then ask about the *simultaneous* impact of all these variables on *mpg*. They would then be led to fit a multiple linear regression model, the result of which is tabulated in Table 3. Many concepts can be explained through this exercise, e.g. *partial regression coefficients*, *partial correlation coefficients*, etc. The analysis of variance tables generated by the computer are useful for explaining additional concepts concerning the various *sums of squares*. Should it be desired, students can be led to discuss the assumptions underlying the multiple linear regression model and the consequences when such assumptions are violated.





